

In-Class Problems — Week 12, Wed

Problem 1. What are each of the following quantities when n independent Bernoulli trials are carried out with probability of success p_s ? (Note: at most you are allowed n trials, not an infinite number of trials)

- (a) The probability of no failures.
- (b) The probability of at least one failure.
- (c) The probability of at most one failure.
- (d) The expected number of failures in n trials.
- (e) The expected number of trials for the first failure.

Problem 2. Each bag of Doritos contains a cool sticker. There are n different kinds of sticker, and I want to collect at least one sticker of each kind. (Assume that sticker kinds in Dorito bags are uniformly random and mutually independent.)

- (a) Suppose that I have already collected k kinds of sticker. What is the expected number of additional bags of Doritos that I must eat to collect one additional kind of sticker?
- (b) What is the expected number of bags of Doritos that I must eat to collect at least one sticker of each kind?

Problem 3. (a) Suppose that I roll a 4-sided die, a 6-sided die, an 8-sided die, a 10-sided die, a 12-sided die, and a 20-sided die. What is the expected number of 6's that come up? (Assume that all of the dice are fair.)

(b) Suppose that I roll n dice that are 6-sided, fair, and mutually independent. What is the expected value of the largest number that comes up?

Hint: You may want to use the “alternative” formula for expectation: $E[M] = \sum_{i=0}^{\infty} \Pr\{M > i\}$. (This formula is valid since the random variables involved are non-negative.)

A Appendix

Definition 3.1. The *expectation* of random variable, R , is:

$$E[R] ::= \sum_{r \in \text{range}(R)} r \cdot \Pr\{R = r\} \quad (1)$$

If R has range \mathbb{N} , then this definition can also be written as

$$E[R] = \sum_{r \in \mathbb{N}} \Pr\{R > r\}$$

Series:

$$\begin{aligned} \sum_{i=0}^{\infty} x^i &= \frac{1}{1-x}, \text{ if } |x| < 1 \\ \sum_{i=0}^n x^i &= \frac{1-x^{n+1}}{1-x} \\ \sum_{i=1}^{\infty} ix^i &= \frac{x}{(1-x)^2} \\ \sum_{i=0}^n i \binom{n}{i} p^i (1-p)^{n-i} &= np \\ H_n &= \sum_{j=1}^n \frac{1}{j} \sim \ln n \end{aligned}$$