

## Problems for Recitation 10

$$1 + z + z^2 + \dots + z^{n-1} = \frac{1 - z^n}{1 - z} \quad (z \neq 1)$$

$$1 + z + z^2 + \dots = \frac{1}{1 - z} \quad (|z| < 1)$$

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n + \frac{1}{2})(n+1)}{3}$$

$$1^3 + 2^3 + 3^3 + \dots + n^3 = \frac{n^2(n+1)^2}{4}$$

**Theorem (Taylor's theorem).** Suppose that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is  $n + 1$  times differentiable on the interval  $[0, x]$ . Then

$$f(x) = f(0) + f'(0)x + \frac{f''(0)x^2}{2!} + \dots + \frac{f^{(n)}(0)x^n}{n!} + \frac{f^{(n+1)}(z)x^{n+1}}{(n+1)!}$$

for some  $z \in [0, x]$ .

# 1 Sums and Approximations

**Problem 1.** Evaluate the following sums.

(a)

$$1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots$$

(b)

$$1 - \frac{1}{2} + \frac{1}{2^2} - \frac{1}{2^3} + \dots$$

(c)

$$1 + 2 + 4 + 8 + \dots + 2^{n-1}$$

(d)

$$\sum_{k=n}^{2n} k^2$$

(e)

$$\sum_{i=0}^n \sum_{j=0}^m 3^{i+j}$$

**Problem 2.** You've seen this neat trick for evaluating a geometric sum:

$$\begin{aligned}S &= 1 + z + z^2 + \dots + z^n \\zS &= z + z^2 + \dots + z^n + z^{n+1} \\S - zS &= 1 - z^{n+1} \\S &= \frac{1 - z^{n+1}}{1 - z}\end{aligned}$$

Use the same approach to find a closed-form expression for this sum:

$$T = 1z + 2z^2 + 3z^3 + \dots + nz^n$$

**Problem 3.** Here is a nasty product:

$$\left(1 + \frac{1}{n^2}\right) \left(1 + \frac{2}{n^2}\right) \left(1 + \frac{3}{n^2}\right) \cdots \left(1 + \frac{n}{n^2}\right)$$

Remarkably, an expression similar to this one comes up in analyzing the distribution of birthdays. Let's make sense of it.

- (a) Give a two-term Taylor approximation for  $e^x$ . (Forget about the error term.)
  
  
  
  
  
  
  
  
  
  
- (b) This is probably the most wide-used approximation in computer science. The fact that  $x$  appears at "ground level" in the approximation and in the exponent of  $e^x$  lets us translate sums into products and vice-versa. Rewrite the product using this approximation.
  
  
  
  
  
  
  
  
  
  
- (c) Now use a standard summation formula to simplify the exponent.
  
  
  
  
  
  
  
  
  
  
- (d) What constant does this approach for large  $n$ ?

**Problem 4.** Let's use Taylor's Theorem to find some approximations for the function  $\sqrt{1+x}$ .

(a) Give a three-term Taylor approximation for  $\sqrt{1+x}$ .

(b) Sketch the function  $\sqrt{1+x}$  and your approximation. How good is the approximation when  $x = 8$ ?

(c) Using this approximation and the fact that  $\sqrt{1+x} = \sqrt{x}\sqrt{1+1/x}$ , give an approximation for  $\sqrt{1+x}$  that is accurate for *large*  $x$

(d) Estimate:

$$\sqrt{1,000,001}$$

to a dozen places beyond the decimal point. You can try to check your answer with a calculator, but you'd better use a good one!