

## Recitation 1: Math Review

February 8, 2007

Elena Grigorescu

## Definitions and Notation

**Problem 1:** Define the following words, phrases and symbols.

1. Set  $A = \{x, y\}$ , subset  $B \subseteq A$ , proper subset  $B \subset A$ , multiset  $\{x, y, y\}$ , power set  $P(A)$ , cardinality  $|A|$ , infinite set, natural numbers  $\mathbb{N} = \{1, 2, 3, \dots\}$ <sup>1</sup>, integers  $\mathbb{Z} = \{\dots, -1, 0, 1, \dots\}$ , empty set  $\emptyset$ , union  $A \cup B$ , intersection  $A \cap B$ , Cartesian product  $A \times B$ , complement  $\bar{A}$ , sequence  $(x, y)$ ,  $k$ -tuple  $(x_1, x_2, \dots, x_k)$ .
2. Function  $f : D \rightarrow R$ , domain  $D$ , range  $R$ , mapping  $\rightarrow$ , one-to-one, onto, bijection (one-to-one, onto).
3. Relation  $R = \{(d_1, r_1), (d_2, r_2), \dots, (d_i, r_i)\}$ , reflexive  $\forall x, xRx$ , symmetric  $\forall x, y, xRy$  iff  $yRx$ , transitive  $\forall x, y, z, xRy \wedge yRz \Rightarrow xRz$ , equivalence (reflexive, symmetric, transitive).
4. Graph  $G = (V, E)$ , degree, path, simple path, cycle, strongly connected.
5. Alphabet (input/output)  $\Sigma = \{a, b, c\}$ , symbols  $a$ , string  $w = baac$ , length  $|w|$ , empty string  $\epsilon$ , substring (consecutive)  $baa$ , concatenation  $w||w$  or  $ww$ , lexicographic ordering  $(\epsilon, 0, 1, 00, 01, 10, 11, 000, \dots)$ <sup>2</sup>, language  $L = \{w_1, w_2, \dots, w_\ell\}$ .
6. Boolean logic  $\{0, 1\}$ , NOT  $\neg p$ , AND  $p \wedge q$ , OR  $p \vee q$ , XOR  $p \oplus q$ , implication  $p \Rightarrow q$ , equality  $p \Leftrightarrow q$ , distributive law  $p \wedge (q \vee r) \Leftrightarrow (p \wedge q) \vee (p \wedge r)$ .
7. Theorem, lemma, corollary, proof, intuition, induction (assumes  $P(n)$ ), strong induction (assumes  $P(0), P(1), \dots, P(n)$ ).
8. (\*) Machine, Automata, language accepted by a machine, language recognized by a machine.

## Proof Techniques

**Problem 2: Set-Theoretic Equivalence:** Recall that in order to prove two sets  $A, B$  are equivalent, one must show that  $A \subseteq B$  and  $B \subseteq A$ . Prove De Morgan's Law that  $\overline{A \cap B} = \bar{A} \cup \bar{B}$ .

**Problem 3: Proof by Contradiction:** 1. If there are 6 people at a party shaking hands, then there must be at least two people who shook hands with the same number of other people.

2. Generalization (Problem 0.12 from Sipser's Text): In any graph with at least 2 nodes there are two nodes of equal degree.

**Problem 4: Proof by Induction (Base Case) (Induction Hypothesis) (Inductive Step):** Problem 0.11 from Sipser's Text.

Find the *error* in the following proof that all horses are the same color.

**Claim:** In any set of  $h$  horses, all horses in the set are the same color.

**Proof:** By induction on  $h$ .

<sup>1</sup>Sipser, pg 4. Zero can also be included in  $\mathbb{N}$ .

<sup>2</sup>Observe the anomaly that 11 precedes 000; length takes precedence.

**Basis:** For  $h = 1$ . In any set containing just one horse, all horses clearly are the same color.

**Inductive Step:** For  $k \geq 1$ , assume that the claim is true for  $h = k$  and prove that it is true for  $h = k + 1$ . Take any set  $H$  of  $k + 1$  horses. we will show that all horses in this set are the same color. Remove one horse from this set to obtain the set  $H_1$  with just  $k$  horses. By the induction hypothesis, all the horses in  $H_1$  are the same color. Now replace the removed horse and remove a different one to obtain the set  $H_2$ . By the same argument, all the horses in  $H_2$  are the same color. Therefore, all the horses in  $H$  must be the same color and the proof is complete.

**Problem 5: Proof by Induction (Base Case) (Induction Hypothesis) (Inductive Step):** Now *correctly* prove the following statement:  $\forall n \in \mathbb{N}, n^3 - n$  is divisible by 6.