1 Linear Programming

1.1 Introduction

Problem description:

- motivate by min-cost flow
- bit of history
- everything is LP
- NP and coNP. P breakthrough.
- general form:
  - **variables**
  - **constraints:** linear equalities and inequalities
  - x **feasible** if satisfies all constraints
  - LP feasible if some feasible x
  - x **optimal** if optimizes objective over feasible x
  - LP is **unbounded** if have feasible x of arbitrary good objective value
  - **lemma:** every lp is infeasible, has opt, or is unbounded
  - (by compactness of R^n and fact that polytopes are closed sets).

Problem formulation:

- canonical form: $\min c^T x, Ax \geq b$
- matrix representation, componentwise $\leq$
- rows a_i of A are **constraints**
- c is **objective**
- any LP has transformation to canonical:
  - max/min objectives same
  - move vars to left, consts to right
  - negate to flip $\leq$ for $\geq$
  - replace $=$ by two $\leq$ and $\geq$ constraints
- standard form: $\min c^T x, Ax = b, x \geq 0$
  - slack variables
  - splitting positive and negative parts $x \to x^+ - x^-$
- $Ax \geq b$ often nicer for theory; $Ax = b$ good for implementations.

point A. 20 minutes.
Some steps towards efficient solution:

- What does answer look like? Can it be represented effectively?
- Easy to verify it is correct?
- Is there a small proof of no answer?
- Can answer, nonanswer be found efficiently?

1.2 Linear Equalities

How solve? First review systems of linear equalities.

- $Ax = b$. when have solution?
- baby case: A is squre matrix with unique solution.
- solve using, eg, Gaussian elimination.
- discuss polynomiality, integer arithmetic later
- equivalent statements:
  - A invertible
  - A^T invertible
  - $\det(A) \neq 0$
  - A has linearly independent rows
  - A has linearly independent columns
  - $Ax = b$ has unique solution for every b
  - $Ax = b$ has unique solution for some b.

What if A isn't square?

- $Ax = b$ has a *witness* for true: give x.
- How about a proof that there is no solution?
- note that "$Ax = b$" means columns of A span b.
- in general, set of points $\{Ax \mid x \in \Re^n\}$ is a **subspace**
- claim: no solution iff for some y, $yA = 0$ but $yb \neq 0$.
- proof: if $Ax = b$, then $yA = 0$ means $yb = yAx = 0$.
- if no $Ax = b$, means columns of A don't span b
- set of points $\{Ax\}$ is subspace not containing b

- find part of b perpendicular to subspace, call it y
- then $yb \neq 0$, but $yA = 0$,
- standard form LP asks for linear combo to, but requires that all coefficients of combo be nonnegative!

Algorithmic?

- Solution means columns of A span b
- Use Gram-Schmidt to reduce A to maximum set of independent columns
- Now maybe more rows (n) than columns (m)?
- But certainly, first m rows of A span first m rows of b
- Solve "square" $Ax = b$ problem
- We know solution is unique if exists
- So must be only solution for all m rows—either works or doesn't.

To talk formally about polynomial size/time, need to talk about size of problems.

- number n has size $\log n$
- rational p/q has size size(p)+size(q)
- size(product) is sum(sizes).
- dimension n vector has size n plus size of number
- $m \times n$ matrix similar: mn plus sizeof numbers
- size (matrix product) at most sum of matrix sizes
- our goal: polynomial time in size of input, measured this way

Claim: if A is $n \times n$ matrix, then $\det(A)$ is poly in size of A

- more precisely, twice the size
- proof by writing determinant as sum of permutation products.
- each product has size n times size of numbers
- $n!$ products
- so size at most size of ($n!$ times product) $\leq n \log n + n \cdot$size(largest entry).

Corollary:

- inverse of matrix is poly size (write in terms of cofactors)
- solution to $Ax = b$ is poly size (by inversion)

1.3 Geometry

Polyhedra

- canonical form: $Ax \geq b$ is an intersection of (finitely many) halfspaces, a **polyhedron**
- standard form: $Ax = b$ is an intersection of hyperplanes (thus a subspace), then $x \geq 0$ intersects in some halfspace. Also a polyhedron, but not full dimensional.
- polyhedron is **bounded** if fits inside some box.
- either formulation defines a **convex** set:
  - if $x, y \in P$, so is $\lambda x + (1 - \lambda)y$ for $\lambda \in 0, 1$.
  - that is, line from x to y stays in P.
- halfspaces define convex sets. Converse also true!
- let C be any convex set, $z \notin C$.
- then there is some a, b such that $ax \geq b$ for $x \in C$, but $az < b$.
- proof by picture. also true in higher dimensions (don't bother proving)
- deduce: every convex set is the intersection of the halfspaces containing it.

1.4 Basic Feasible Solutions

Again, let's start by thinking about structure of optimal solution.

- Can optimum be in "middle" of polyhedron?
- Not really: if can move in all directions, can move to improve opt.

Where can optimum be? At "corners."

- "vertex" is point that is not a convex combination of two others
- "extreme point" is point that is *unique* optimum in some direction

Basic solutions:

- A constraint $ax \leq b$ or $ax = b$ is *tight* or *active* if $ax = b$
- for n-dim LP, point is *basic* if (i) all equality constraints are tight and (ii) n linearly independent constraints are tight.
- in other words, x is at intersection of boundaries of n linearly independent constraints

- note x is therefore the unique intersection of these boundaries.
- a *basic feasible solution* is a solution that is basic and satisfies all constraints.

In fact, vertex, extreme point, bfs are *equivalent*.

- Proof left somewhat to reader.

Any standard form lp $\min cx,\ Ax = b,\ x \geq 0$ with opt has one at a BFS.

- Suppose opt x is not at BFS
- Then less than n tight constraints
- So at least one degree of freedom
- i.e, there is a (linear) subspace on which all those constraints are tight.
- In particular, some line through x for which all these constraints are tight.
- Write as $x + \epsilon d$ for some vector direction d
- Since x is feasible and other constraints *not* tight, $x + \epsilon d$ is feasible for small enough ϵ.
- Consider moving along line. Objective value is $cx + \epsilon cd$.
- So for either positive or negative ϵ, objective is *nonincreasing*, i.e. doesn't get worse.
- Since started at opt, must be no change at all—i.e., $cd = 0$.
- So can move in *either* direction.
- In at least one direction, some x_i is decreasing.
- Keep going till new constraint becomes tight (some $x_i = 0$).
- Argument can be repeated until n tight constraints, i.e. bfs
- Conclude: every standard form LP with an optimum has one at a bfs.
- Note convenience of using standard form: ensures bounded, so can reach bfs
- canonical form has oddities: e.g. $\max y \mid y \leq 1$.
- but any *bounded, feasible* LP has BFS optimum

Other characterizations of corner:

- "vertex" is point that is not a convex combination of two others
- "extreme point" is point that is *unique* optimum in some direction

- Previous proof shows extreme point is BFS (because if cannot move to any other opt, must have n tight constraints).
- Also shows BFS is vertex:
  - if point is convex combo (not vertex), consider line through it
  - all points on it feasible
  - so don't have n tight constraints
  - conversely, if less than n tight constraints, they define feasible subspace containing line through point
  - so point is convex combo of points on line.
- To show BFS is extreme point, show point is unique opt for objective that is sum of normals to tight constraints.

Yields first algorithm for LP: try all bfs.

- How many are there?
- just choose n tight constraints out of m, check feasibility and objective
- Upper bound $\binom{m}{n}$

Also shows output is polynomial size:

- Let A' and correspoinding b' be n tight constraints (rows) at opt
- Then opt is (unique) solution to $A'x = b'$
- We saw last time that such an inverse is represented in polynomial size in input

(So, at least *weakly* polynomial algorithms seem possible)
Corollary:

- Actually showed, if x feasible, exists BFS with no worse objective.
- Note that in canconical form, might not have opt at vertex (optimize x_1 over (x_1, x_2) such that $0 \leq x_1 \leq 1$).
- But this only happens if LP is unbounded
- In particular, if opt is *unique*, it is a bfs.

OK, this is an exponential method for finding the optimum. Maybe we can do better if we just try to verify the optimum. Let's look for a way to prove that a given solution x is optimal.
Quest for nonexponential algorithm: start at an easier place: how decide if a solution is optimal?

- decision version of LP: is there a solution with opt$> k$?
- this is in NP, since can exhibit a solution (we showed poly size output)
- is it in coNP? Ie, can we prove there is no solution with opt$> k$? (this would give an optimality test)

2 Duality

What about optimality?

- Intro *duality*, strongest result of LP
- give proof of optimality
- gives max-flow mincut, prices for mincost flow, game theory, lots other stuff.

Motivation: find a **lower** bound on $z = \min\{cx \mid Ax = b, x \geq 0\}$.

- Standard approach: try adding up combos of existing equations
- try multiplying $a_i x = b_i$ by some y_i. Get $yAx = yb$
- If find y s.t. $yA = c$, then $yb = yAx = cx$ and we know opt (we inverted $Ax = b$)
- looser: if require $yA \leq c$, then $yb = yAx \leq cx$ is lower bound since $x_j \geq 0$
- so to get best lower bound, want to solve $w = \max\{yb \mid yA \leq c\}$.
- this is a new linear program, *dual* of original.
- just saw that dual is less than primal (weak duality)

Note: dual of dual is primal:

$$\begin{aligned}
\max\{yb : yA \leq c\} &= \max\{by \mid A^T y \leq c\} \\
&= -\min\{-by \mid A^T y + Is = c, s \geq 0\} \\
&= -\min\{-by^+ + by^- \mid A^T y + (-A^T)y^- + Is = c, y^+, y^-, s \geq 0\} \\
&= -\max\{cz \mid zA^T \leq -b, z(-A^T) \leq -b, Iz \leq 0\} \\
&= \min\{cx \mid Ax = b, x \geq 0\} \quad (x = -z)
\end{aligned}$$

Weak duality: if P (min, opt z) and D (max, opt w) feasible, $z \geq w$

- $w = yb$ and $z = cx$ for some primal/dual feasible y, x
- x primal feasible ($Ax = b, x \geq 0$)
- y dual feasible ($yA \leq c$)
- then $yb = yAx \leq cx$

Note corollary:

- (restatement:) if P, D both feasible, then both bounded.
- if P feasible and unbounded, D not feasible
- if P feasible, D either infeasible or bounded

- in fact, only 4 possibilities. both feasible, both infeasible, or one infeasible and one unbounded.
- **notation:** P unbounded means D infeasible; write solution $-\infty$. D unbounded means P infeasilbe, write solution ∞.

3 Strong Duality

Strong duality: if P or D is feasible then $z = w$

- includes D infeasible via $w = -\infty$)

Proof by picture:

- $\min\{yb \mid yA \geq c\}$ (note: **flipped sign**)
- suppose b points straight up.
- imagine ball that falls down (minimize height)
- stops at opt y (no local minima)
- stops because in physical equilibrium
- equilibrium exterted by forces normal to "floors"
- that is, aligned with the A_i (columns)
- but those floors need to cancel "gravity" $-b$
- thus $b = \sum A_i x_i$ for some **nonnegative** force coeffs x_i.
- in other words, x feasible for $\min\{cx \mid Ax = b, x \geq 0\}$
- also, only floors touching ball can exert any force on it
- thus, $x_i = 0$ if $yA_i > c_i$
- that is, $(c_i - yA_i)x_i = 0$
- thus, $cx = \sum(yA_i)x_i = yb$
- so x is dual optimal.

Let's formalize.

- Consider optimum y
- WLOG, ignore all loose constraints (won't need them)
- And if any are redundant, drop them
- So at most n tight constraints remain

- and all linearly independent.
- and since those constraints are tight, $yA = c$

Claim: Exists x, $Ax = b$

- Suppose not? Then "duality" for linear equalities proves exists z, $zA = 0$ but $zb <> 0$.
- WLOG $zb < 0$ (else negate it)
- So consider $y + z$.
- $A(y + z) = Ay + Az = Ay$, so feasible
- $b(y + z) = by + bz < by$, so better than opt! Contra.

Claim: $yb = cx$

- Just said $Ax = b$ in dual
- In primal, all (remaining) constraints are tight, so $yA = c$
- So $yb = yAx = cx$

Claim: $x \geq 0$

- Suppose not.
- Then some $x_i < 0$
- Let $c' = c + e_i$
- In other words, moving i^{th} constraint $yA_i \geq c_i$ "upwards"
- Consider solution to $y'A = c'$
- Exists solution (since A is full rank)
- And $c' \geq c$, so $y'A = c'$ is *feasible* for original constraints $yA \geq c$
- Value of objective is $y'b = y'Ax = c'x$
  - We assumed $x_i < 0$, and *increased* c_i
  - So $c'x < cx$
  - So got better value than opt. Contradiction!
- Intuition: x_i is telling us how much opt will change if we "tighten" i^{th} constraint

Neat corollary: Feasibility or optimality: which harder?

- given optimizer, can check feasiblity by optimizing arbitrary func.
- Given feasibility algorithm, can optimize by combining primal and dual.

Interesting note: knowing dual solution may be useless for finding optimum (more formally: if your alg runs in time T to find primal solution given dual, can adapt to alg that runs in time $O(T)$ to solve primal without dual).

3.1 Rules for duals

General dual formulation:

- primal is

$$\begin{aligned} z &= \min c_1x_1 + c_2x_2 + c_3x_3 \\ A_{11}x_1 + A_{12}x_2 + A_{13}x_3 &= b_1 \\ A_{21}x_1 + A_{22}x_2 + A_{23}x_3 &\geq b_2 \\ A_{31}x_1 + A_{32}x_2 + A_{33}x_3 &\leq b_3 \\ x_1 &\geq 0 \\ x_2 &\leq 0 \\ x_3 & \quad UIS \end{aligned}$$

  (UIS emphasizes unrestricted in sign)

- means dual is

$$\begin{aligned} w &= \max y_1b_1 + y_2b_2 + y_3b_3 \\ y_1A_{11} + y_2A_{21} + y_3A_{31} &\leq c_1 \\ y_1A_{12} + y_2A_{22} + y_3A_{32} &\geq c_2 \\ y_1A_{13} + y_2A_{23} + y_3A_{33} &= c_3 \\ y_1 & \quad UIS \\ y_2 &\geq 0 \\ y_3 &\leq 0 \end{aligned}$$

- In general, variable corresponds to constraint (and vice versa):

PRIMAL	minimize	maximize	DUAL
constraints	$\geq b_i$ $\leq b_i$ $= b_i$	≥ 0 ≤ 0 free	variables
variables	≥ 0 ≥ 0 free	$\leq c_j$ $\leq c_j$ $= c_j$	constraints

Derivation:

- remember lower bounding plan: use $yb = yAx \leq cx$ relation.
- If constraint is in "natural" direction, dual variable is positive.
- We saw A_{11} and x_1 case. $x_1 \geq 0$ ensured $yAx_1 \leq c_1x_1$ for **any** y

- If some $x_2 \leq 0$ constraint, we want $yA_{12} \geq c_2$ to maintain rule that $y_1 A_{12} x_2 \leq c_2 x_2$
- If x_3 unconstrained, we are only safe if $yA_{13} = c_3$.
- if instead have $A_{21}x_1 \geq b_2$, any old y won't do for lower bound via $c_1 x_1 \geq y_2 A_{21} x_1 \geq y_2 b_2$. Only works if $y_2 \geq 0$.
- and so on (good exercise).
- This gives weak duality derivation. Easiest way to derive strong duality is to transform to standard form, take dual and map back to original problem dual (also good exercise).

Note: tighter the primal, looser the dual

- (equality constraint leads to unrestricted var)
- adding primal constraints creates a new dual variable: more dual flexibility

3.2 Shortest Paths

A dual example:

- shortest path is a dual (max) problem:

$$\begin{aligned} w &= \max d_t - d_s \\ d_j - d_i &\leq c_{ij} \end{aligned}$$

- constraints matrix A has ij rows, i columns, ± 1 entries (draw)
- what is primal? unconstrained vars, give equality constraints, dual upper bounds mean vars must be positive.

$$\begin{aligned} z &= \min \sum y_{ij} c_{ij} \\ y_{ij} &\geq 0 \end{aligned}$$

thus

$$\sum_j y_{ji} - y_{ij} = 1(i = s), -1(i = t), 0 \text{ ow}$$

It's the minimum cost to send one unit of flow from s to t!

4 Complementary Slackness

Leads to another idea: *complementary slackness*:

- given feasible solutions x and y, $cx - yb \geq 0$ is *duality gap.*
- optimal iff gap 0 (good way to measure "how far off"
- Go back to original primal and dual forms
- rewrite dual: $yA + s = c$ for some $s \geq 0$ (that is, $s_j = c_j - yA_j$)
- The following are equivalent for feasible x, y:
  - x and y are optimal
  - $sx = 0$
  - $x_j s_j = 0$ for all j
  - $s_j > 0$ implies $x_j = 0$
- We saw this in duality analysis: only tight constraints "push" on opt, giving nonzero dual variables.
- proof:
  - $cx = by$ iff $(yA + s)x = y(Ax)$, iff $sx = 0$
  - if $sx = 0$, then since $s, x \geq 0$ have $s_j x_j = 0$ (converse easy)
  - so $s_j > 0$ forces $x_j = 0$ (converse easy)
- basic idea: opt cannot have a variable x_j and corresponding dual constraint s_j slack at same time: one must be tight.
- Generalize to arbitrary form LPs: feasible points optimal if:

$$\begin{aligned} y_i(a_i x - b_i) &= 0 \forall i \\ (c_j - yA_j)x_j &= 0 \forall j \end{aligned}$$

- proof:
  - note in definition of dual, $\geq$ constraint in primal (min) corresponds to nonnegative y_i.
  - thus, feasiblity means $y_i(a_i x - b_i) \geq 0$.
  - similarly, $\leq$ constraint in dual corresponds to nonnegative x_j in primal
  - so feasibility means $(c_j - yA_j)x_j \geq 0$.

- Also,

$$\begin{aligned}\sum y_i(a_i x - b_i) + (c_j - yA_j)x_j &= yAx - yb + cx - yAx \\ &= cx - yb \\ &= 0\end{aligned}$$

  at opt. But since just argued all terms are nonnegative, all must be 0

- conversely, if all are 0, then $cx = by$, so we are optimal

Let's take some duals.
Max-Flow min-cut theorem:

- modify to circulation to simplify
- primal problem: create infinite capacity (t, s) arc

$$\begin{aligned}P &= \max \sum_w x_{ts} \\ \sum_w x_{vw} - x_{wv} &= 0 \\ x_{vw} &\leq u_{vw} \\ x_{vw} &\geq 0\end{aligned}$$

- dual problem: vars z_v dual to balance constraints, y_{vw} dual to capacity constraints.

$$\begin{aligned}D &= \min \sum_{vw} y_{vw} u_{vw} \\ y_{vw} &\geq 0 \\ z_v - z_w + y_{vw} &\geq 0 \\ z_t - z_s + y_{ts} &\geq 1\end{aligned}$$

- Think of y_{vw} as "lengths"
- note $y_{ts} = 0$ since otherwise dual infinite. so $z_t - z_s \geq 1$.
- rewrite as $z_w \leq z_v + y_{vw}$.
- deduce y_{vw} are edge lengths, z_v are "distances"
- In particular, can substract z_s from everything without changing feasibility (subs cancel)
- Now z_v is upper bound on distance from source to v.
- So, are trying to maximize source-sink distance

- Good justification for shortest aug path, blocking flows

- sanity check: mincut: assign length 1 to each mincut edge
- unfortunately, might have noninteger dual optimum.
- let $S = v \mid z_v < 1$ (so $s \in S, t \notin S$)
- use complementary slackness:
  - if (v, w) *leaves* S, then $y_{vw} \geq z_w - z_v > 0$, so $x_{vw} = u_{vw}$, (tight) i.e. (v, w) saturated.
  - if (v, w) *enters* S, then $z_v > z_w$. Also know $y_{vw} \geq 0$; add equations and get $z_v + y_{vw} > z_w$ i.e. slack.
  - so $x_{wv} = 0$
  - in other words: all leaving edges saturated, all coming edges empty.
- now just observe that value of flow equals value crossing cut equals value of cut.

Min cost circulation: change the objective function associated with max-flow.

- primal:

$$\begin{aligned} z &= \min \sum c_{vw} x_{vw} \\ \sum_w x_{vw} - x_{wv} &= 0 \\ x_{vw} &\leq u_{vw} \\ x_{vw} &\geq 0 \end{aligned}$$

- as before, dual: variable y_{vw} for capacity constraint on f_{vw}, z_v for balance.
- Change to primal min problem flips sign constraint on y_{vw}
- What does change in primal objective mean for dual? Different constraint bounds!

$$\begin{aligned} & \max \sum y_{vw} u_{vw} \\ z_v - z_w + y_{vw} &\leq c_{vw} \\ y_{vw} &\leq 0 \\ z_v & \quad \text{UIS} \end{aligned}$$

- rewrite dual: $p_v = -z_v$

$$\begin{aligned} & \max \sum y_{vw} u_{vw} \\ y_{vw} &\leq 0 \\ y_{vw} &\leq c_{vw} + p_v - p_w = c_{vw}^{(p)} \end{aligned}$$

- Note: $y_{vw} \leq 0$ says the objective function is the sum of the **negative parts** of the reduced costs (positive ones get truncated to 0)
- Note: optimum ≤ 0 since of course can set $y = 0$. Since since zero circulation is primal feasible.
- complementary slackness.
  - Suppose $f_{vw} < u_{vw}$.
  - Then dual variable $y_{vw} = 0$
  - So $c_{ij}^{(p)} \geq 0$
  - Thus $c_{ij}^{(p)} < 0$ implies $f_{ij} = u_{ij}$
  - that is, all negative reduced cost arcs saturated.
  - on the other hand, suppose $c_{ij}^{(p)} > 0$
  - then constraint on z_{ij} is slack
  - so $f_{ij} = 0$
  - that is, all positive reduced arcs are empty.

5 Algorithms

5.1 Simplex

vertices in standard form/bases:

- Without loss of generality make A have full row rank (define):
  - find basis in rows of A, say $a_1, \ldots, a_k$
  - any other a_ℓ is linear combo of those.
  - so $a_\ell x = \sum \lambda_i a_i x$
  - so better have $b_l = \sum \lambda_i a_i$ if any solution.
  - if so, anything feasible for $a_1, \ldots, a_\ell$ feasible for all.
- m constraints $Ax = b$ all tight/active
- given this, need $n - m$ of the $x_i \geq 0$ constraints
- also, need them to form a basis with the a_i.
- **write matrix** of tight constraints, first m rows then identity matrix
- need linearly independent rows
- equiv, need linearly independent columns
- but columns are linearly independent iff m columns of A including all corresp to nonzero x are linearly independent

- gives other way to define a vertex: x is vertex if
  - $Ax = b$
  - m linearly independent columns of A include all $x_j \neq 0$

  This set of m columns is called a *basis*.
- x_j of columns called *basic* set B, others *nonbasic* set N
- given bases, can compute x:
  - A_B is basis columns, $m \times m$ and full rank.
  - solve $A_B x_B = b$, set other $x_N = 0$.
  - note can have many bases for same vertex (choice of 0 x_j)

Summary: x is vertex of P if for some basis B,

- $x_N = 0$
- A_B nonsingular
- $A_B^{-1} b \geq 0$

Simplex method:

- start with a basic feasible soluion
- try to improve it
- rewrite LP: $\min c_B x_B + c_N x_N$, $A_B x_B + A_N x_N = b$, $x \geq 0$
- B is basis for bfs
- since $A_B x_B = b - A_N x_N$, so $x_B = A_B^{-1}(b - A_N x_N)$, know that

$$\begin{aligned} cx &= c_B x_B + c_N x_N \\ &= c_B A_B^{-1}(b - A_N x_N) + c_N x_N \\ &= c_B A_B^{-1} b + (c_N - c_B A_B^{-1} A_N) x_N \end{aligned}$$

- *reduced cost* $\tilde{c}_N = c_N - c_B A_B^{-1} A_N$
- if no $\tilde{c}_j < 0$, then increasing any x_j increases cost (may violate feasiblity for x_B, but who cares?), so are at optimum!
- if some $\tilde{c}_j < 0$, can increase x_j to decrease cost
- but since x_B is func of x_N, will have to stop when x_B hits a constraint.
- this happens when some x_i, $i \in B$ hits 0.

- we bring j into basis, take i out of basis.
- we've moved to an *adjacent* basis.
- called a *pivot*
- **show picture**

Notes:

- Need initial vertex. How find?
- maybe some $x_i \in B$ already 0, so can't increase x_j, just pivot to same obj function.
- could lead to cycle in pivoting, infinite loop.
- can prove exist noncycling pivots (eg, lexicographically first j and i)
- no known pivot better than exponential time
- note traverse path of edges over polytope. Unknown what shortest such path is
- Hirsh conjecture: path of $m - d$ pivots exists.
- even if true, simplex might be bad because path might not be monotone in objective function.
- certain recent work has shown $n^{\log n}$ bound on path length

5.2 Simplex and Duality

- defined *reduced costs* of nonbasic vars N by

$$\tilde{c}_N = c_N - c_B A_B^{-1} A_N$$

  and argued that when all $\tilde{c}_N \geq 0$, had optimum.
- Define $y = c_B A_B^{-1}$ (so of course $c_B = y A_B$)
- nonegative reduced costs means $c_N \geq y A_N$
- put together, see $yA \leq c$ so y is dual feasible
- but, $yb = c_B A_B^{-1} b = c_B x_B = cx$ (since $x_N = 0$)
- so y is dual optimum.
- more generally, y measures duality gap for current solution!
- another way to prove duality theorem: prove there is a terminating (non cycling) simplex algorithm.

5.3 Polynomial Time Bounds

We know a lot about structure. And we've seen how to verify optimality in polynomial time. Now turn to question: can we solve in polynomial time?
Yes, sort of (Khachiyan 1979):

- polynomial algorithms exist
- strongly polynomial unknown.

Claim: all vertices of LP have polynomial size.

- vertex is bfs
- bfs is intersection of n constraints $A_B x = b$
- invert matrix.

Now can prove that feasible alg can optimize a different way:

- use binary search on value z of optimum
- add constraint $cx \leq z$
- know opt vertex has poly number of bits
- so binary search takes poly (not logarithmic!) time
- not as elegant as other way, but one big advantage: feasiblity test over basically same polytope as before. Might have fast feasible test for this case.

6 Ellipsoid

Lion hunting in the desert.

- bolzano wierstrauss theorem—proves certain sequence has a subsequence with a limit by repeated subdividing of intervals to get a point in the

Define an ellipsoid

- generalizes ellipse
- write some $D = BB^T$ "radius"
- *center* z
- point set $\{(x-z)^T D^{-1}(x-z) \leq 1\}$
- note this is just a basis change of the unit sphere $x^2 \leq 1$.
- under transform $x \to Bx + z$

Outline of algorithm:

- goal: find a feasible point for $P = \{Ax \leq b\}$
- start with ellipse containing P, center z
- check if $z \in P$
- if not, use separating hyperplane to get 1/2 of ellipse containing P
- find a smaller ellipse containing this 1/2 of original ellipse
- until center of ellipse is in P.

Consider sphere case, separating hyperplane $x_1 = 0$

- try center at $(a, 0, 0, \ldots)$
- Draw picture to see constraints
- requirements:
  - $d_1^{-1}(x_1 - a)^2 + \sum_{i>1} d_i^{-1} x_i^2 \leq 1$
  - constraint at $(1, 0, 0)$: $d_1^{-1}(x - a)^2 = 1$ so $d_1 = (1 - a)^2$
  - constraint at $(0, 1, 0)$: $a^2/(1-a)^2 + d_2^{-1} = 1$ so $d_2^{-1} = 1 - a^2/(1-a)^2 \approx 1 - a^2$
- What is volume? about $(1 - a)/(1 - a^2)^{n/2}$
- set a about $1/n$, get $(1 - 1/n)$ volume ratio.

Shrinking Lemma:

- Let $E = (z, D)$ define an n-dimensional ellipsoid
- consider separating hyperplane $ax \leq az$
- Define $E' = (z', D')$ ellipsoid:

$$\begin{aligned} z' &= z - \frac{1}{n+1} \frac{Da^T}{\sqrt{aDa^T}} \\ D' &= \frac{n^2}{n^2 - 1}(D - \frac{2}{n+1} \frac{Da^T aD}{aDa^T}) \end{aligned}$$

- then

$$\begin{aligned} E \cap \{x \mid ax \leq ez\} &\subseteq E' \\ \mathrm{vol}(E') &\leq e^{1/(2n+1)} \mathrm{vol}(E) \end{aligned}$$

- for proof, first show works with $D = I$ and $z = 0$. new ellipse:

$$\begin{aligned} z' &= -\frac{1}{n+1} \\ D' &= \frac{n^2}{n^2-1}(I - \frac{2}{n+1} I_{11}) \end{aligned}$$

  and volume ratio easy to compute directly.

- for general case, transform to coordinates where $D = I$ (using new basis B), get new ellipse, transform back to old coordinates, get (z', D') (note transformation don't affect volume *ratios*.

So ellipsoid shrinks. Now prove 2 things:

- needn't start infinitely large
- can't get infinitely small

Starting size:

- recall bounds on size of vertices (polynomial)
- so coords of vertices are exponential but no larger
- so can start with sphere with radius exceeding this exponential bound
- this only uses polynomial values in D matrix.
- if unbounded, no vertices of P, will get vertex of box.

Ending size:

- convenient to assume that polytope full dimensional
- if so, it has $n+1$ affinely indpendent vertices
- all the vertices have poly size coordinates
- so they contain a box whose volume is a poly-size number (computable as determinant of vertex coordinates)

Put together:

- starting volume $2^{n^{O(1)}}$
- ending volume $2^{-n^{O(1)}}$
- each iteration reduces volume by $e^{1/(2n+1)}$ factor
- so $2n+1$ iters reduce by e
- so $n^O(1)$ reduce by $e^{n^{O(1)}}$

- at which point, ellipse doesn't contain P, contra
- must have hit a point in P before.

Justifying full dimensional:

- take $\{Ax \leq b\}$, replace with $P' = \{Ax \leq b + \epsilon\}$ for tiny ϵ
- any point of P is an interior of P', so P' full dimensional (only have interior for full dimensional objects)
- P empty iff P' is (because ϵ so small)
- can "round" a point of P' to P.

Infinite precision:

- built a new ellipsoid each time.
- maybe its bits got big?
- no.

6.1 Separation vs Optimization

Notice in ellipsoid, were only using one constraint at a time.

- didn't matter how many there were.
- didn't need to see all of them at once.
- just needed each to be represented in polynomial size.
- so ellipsoid works, even if huge number of constraints, so long as have *separation oracle:* given point not in P, find separating hyperplane.
- of course, feasibility is same as optimize, so can optimize with sep oracle too.
- this is on a polytope by polytope basis. If can separate a particular polytope, can optimize over that polytope.

This is very useful in many applications. e.g. network design.

7 Interior Point

Ellipsoid has problems in practice ($O(n^6)$ for one). So people developed a different approach that has been extremely successful.
What goes wrong with simplex?

- follows edges of polytope

- complex stucture there, run into walls, etc
- interior point algorithms stay away from the walls, where structure simpler.
- Karmarkar did the first one (1984); we'll descuss one by Ye

7.1 Potential Reduction

Potential function:

- Idea: use a (nonlinear) potential function that is minimized at opt but also enforces feasibility
- use gradient descent to optimize the potential function.
- Recall standard primal $\{Ax = b, x \geq 0\}$ and dual $yA + s = c, s \geq 0$.
- duality gap sx
- Use *logarithmic barrier function*

  $$G(x, s) = q \ln xs - \sum \ln x_j - \sum \ln s_j$$

  and try to minimize it (pick q in a minute)
- first term forces duality gap to get small
- second and third enforce positivity
- note barrier prevents from ever hitting optimum, but as discussed above ok to just get close.

Choose q so first term dominates, guarantees good G is good xs

- $G(x, s)$ small should mean xs small
- xs large should mean $G(x, s)$ large
- write $G = \ln(xs)^q / \prod x_j s_j$
- $xs > x_j s_j$, so $(xs)^n > \prod x_j s_j$. So taking $q > n$ makes top term dominate, $G > \ln xs$

How minimize potential function? Gradient descent.

- have current (x, s) point.
- take linear approx to potential function around (x, s)
- move to where linear approx smaller $(-\nabla_x G)$
- deduce potential also went down.

- crucial: can only move as far as linear approximation accurate

Firs wants big q, second small q. Compromise at $n + \sqrt{n}$, gives $O(L\sqrt{n})$ iterations.
Must stay feasible:

- Have gradient $g = \nabla_x G$
- since potential not minimized, have reasonably large gradient, so a small step will improve potential a lot. **picture**
- want to move in direction of G, but want to stay feasilbe
- project G onto nullspace(A) to get d
- then $A(x + d) = Ax = b$
- also, for sufficiently small step, $x \geq 0$
- potential reduction proportional to length of d
- problem if d too small
- In that case, move s (actually y) by $g - d$ which will be big.
- so can either take big primal or big dual step
- why works? Well, d (perpendicular to A) has $Ad = 0$, so good primal move.
- converseley, part spanned by A has $g - d = wA$,
- so can choose $y' = y + w$ and get $s' = c - Ay' = c - Ay - (g - d) = s - (g - d)$.
- note $dG/dx_j = s_j/(xs) - 1/x_j$
- and $dG/ds_j = x_j/(xs) - 1/s_j = (x_j/s_j)dG/dx_j \approx dG/dx_j$