Cauchy's rigidity theorem [Cauchy 1813; Steinitz 1934]

6.885
L11.4
Oct. 18, 2004

If two convex polyhedra are combinatorially equivalent (same graph structure) and corresponding faces are congruent, then polyhedra are congruent

Proof by contradiction: Consider counterexample P, P'

- for each vertex pair v, v':
  slice with ε-sphere centered at vertex (of polyhedron)
  $\Rightarrow$ spherical polygon, edge lengths = face angles, angles = dihedral angles
- faces congruent $\Rightarrow$ edge lengths match in V vs. V'
- P, P' incongruent $\Rightarrow$ angles don't match for some v, v' pair
- for such a pair v, v':
  - label vertex of v's spherical polygon + if larger angle in V, − if smaller angle in V ($\emptyset$ otherwise)
  - can't be all + (or $\emptyset$) or all − (or $\emptyset$):

    Cauchy arm lemma (plane or sphere): opening all angles of convex open chain increases distance between endpoints

    $\Rightarrow$ edge length not preserved. (would be)



  $\Rightarrow \geq 2$ alternations: $+ + \ldots + - - \ldots -$
  - can't be just 2, by arm lemma:



  $\Rightarrow \geq 4$ alternations: $+ - + -$
  $\Rightarrow$ # alternations $\geq 4V$ in subgraph of +/− edges.
- +/− labels extend to edges of polyhedron
- alternation at vertex $\overset{1\text{-}1}{\Leftrightarrow}$ alternation in incident face
- $\leq 2k$ alternations in face of degree $2k$ or $2k+1$

  $\Rightarrow 4V \leq$ # alternations $\leq 2f_3 + 4f_4 + 4f_5 + 6f_6 + 6f_7 + \cdots$
- $E = \frac{1}{2}(3f_3 + 4f_4 + 5f_5 + 6f_6 + 7f_7 + \cdots)$
- $V = 2 + E - F = 2 + \frac{1}{2}(f_3 + 2f_4 + 3f_5 + 4f_6 + 5f_7 + \cdots)$

$\Rightarrow 4V = 8 + 2f_3 + 4f_4 + 6f_5 + 8f_6 + 10f_7 + \cdots$ $\longrightarrow$ ⊠ □