

Optimization problem: (combinatorial)

- goal: instance \rightarrow solution with min/max cost
- set of instances
- for each instance:
 - set of (valid/feasible) solutions
 - nonnegative cost of each solution (\mathbb{R} or \mathbb{Z})
- objective: min or max

OPT(x) = min/max possible cost for instance x
(sometimes also the solution itself)

NP optimization problem:

- solutions have polynomial length
- instances & their solutions can be recognized $\in P$
- cost function $\in P$

\Rightarrow decision problem $\in NP$

\hookrightarrow min: is $OPT(x) \leq q$? ($\geq \in coNP$)

max: is $OPT(x) \geq q$? ($\leq \in coNP$)

NPO = $\{NP \text{ optimization problems}\}$

Approximation: ALG is a c -approximation if $\forall x$:
- min: $\frac{\text{cost}(\text{ALG}(x))}{\text{cost}(\text{OPT}(x))} \leq c$ ($c \geq 1$) \downarrow instance

- max: $\frac{\text{cost}(\text{OPT}(x))}{\text{cost}(\text{ALG}(x))} \leq c$ ($c \geq 1$) e.g. 2

OR: $\frac{\text{cost}(\text{ALG}(x))}{\text{cost}(\text{OPT}(x))} \geq c$ ($c \leq 1$) e.g. $\frac{1}{2}$

- usually: ALG should be polynomial time

PTAS (Polynomial-Time Approximation Scheme)

= algorithm with additional input $\varepsilon > 0$

- solution is $(1+\varepsilon)$ -approximation

- polynomial time for every fixed $\varepsilon > 0$

- e.g. $n^{2^{1/\varepsilon}}$ OK (tighter notions later)

PTAS = $\{ \text{NP optimization problems having PTAS} \}$
= $\bigcap_{c \geq 1} c\text{-APX}$

F-APX = $\{ \text{NP optimization problems having poly-time } f(n)\text{-approximation algorithm for some } f \in F \}$

APX = $O(1)$ -APX

= MAX SNP in older literature

Log-APX = $O(\lg n)$ -APX

Poly-APX = $n^{O(1)}$ -APX

- $P \neq NP \Rightarrow \text{PTAS} \subsetneq \text{APX} \subsetneq \text{Log-APX} \subsetneq \text{Poly-APX}$ etc.

Typical approximation factors: (graph problems)

- $1+\epsilon$ (PTAS)

- lots of problems on planar/H-minor-free graphs
e.g. H-minor-free dominating set ↷
choose min. # vertices adjacent to unchosen vertices
& in Euclidean plane e.g. TSP,
Steiner tree, rectilinear Steiner tree [L9]

- $\Theta(1)$ (APX-complete)

- lots e.g. TSP, Steiner tree
- max. coverage: choose k vertices from left side of bipartite graph adjacent to max. # vertices ←

- $\Theta(\log^* n)$

- asymmetric k -center: given asymmetric metric, choose k vertices to min. max distance $v \rightarrow$ nearest chosen

- $\Theta(\log n)$

- set cover & dominating set
↳ dominating set from left side of bipartite graph ←
- max. unique coverage (exactly 1 left adjacent to right)

"dual"

- $\Theta(\log^2 n)$

- group Steiner tree: given graph & k groups of vertices, choose min. # vertices inducing connected subgraph & containing at least 1 vertex in each group

- $\Omega(\log^2 n) \cap O(n^\epsilon)$ (OPEN)

- directed Steiner tree: given graph, k terminal vertices, & root vertex, choose min. # vertices inducing root-to-terminal path for each terminal

- $\Omega(2^{\log^{1-\epsilon} n}) \cap O(n^c)$ (OPEN)

$c = \frac{1}{3}$ → - label cover (MinRep & MaxRep) [future lecture]

$c = \frac{4}{5} + \epsilon$ → - directed Steiner forest: given $s_i \rightarrow t_i$ pairs, choose min. # vertices inducing such paths

- $\Omega(n^{1-\epsilon}) \cap \tilde{O}(n)$

↑ polylog factors

- chromatic number: min k such that k -colorable

- independent set \equiv clique (complement graph)

Approximation preserving reductions: $A \rightarrow B$

(see Crescenzi - CCC 1997)

A instance x

\xrightarrow{f} B instance $x' = f(x)$

A solution $y = g(x, y')$ to x \xleftarrow{g} B solution y' to x'

PTAS-reduction: $\forall \epsilon > 0 \exists \delta = \delta(\epsilon) > 0$ such that

y' is $(1 + \delta(\epsilon))$ -approximation to B
 $\Rightarrow y = g(x, y')$ is $(1 + \epsilon)$ -approximation to A

[Crescenzi & Trevisan 1994]

- f & g can depend on ϵ too (else "P-reduction")
- $B \in \text{PTAS} \Rightarrow A \in \text{PTAS}$ (chain algs. together)
- $A \notin \text{PTAS} \Rightarrow B \notin \text{PTAS}$
- ditto for APX
- careful: $A \in \text{PTAS} \not\Rightarrow B \in \text{PTAS}$
- if $\delta(\emptyset) = \emptyset$ also works then also NP reduction
- reductions chain: $A \rightarrow B \rightarrow C$

AP-reduction: $\delta(\epsilon) = O(\epsilon)$

- $B \in O(f)$ -APX $\Rightarrow A \in O(f)$ -APX

[Crescenzi, Kann, Silvestri, Trevisan 1995]

Strict reduction: $\delta(\epsilon) = \epsilon$

[Orponen & Mannila 1987]

APX-hard = \exists PTAS-reduction from any problem \in APX
 - \notin PTAS if $P \neq NP$

$O(f)$ -APX-hard = \exists AP-reduction from any problem $\in O(f)$ -APX
 (other definitions possible)
 - $\notin O(f)$ -APX if $P \neq NP$

L-reduction: $OPT_B(x') = O(\overset{\rightarrow \leq \alpha}{OPT_A(x)})$
 & $|cost_A(y) - OPT_A(x)| = O(\overset{\leftarrow \leq \beta}{|cost_B(y') - OPT_B(x')|})$
 [Papadimitriou & Yannakakis - JCSS 1991]

\Rightarrow PTAS-reduction

- for minimization problems:

\Rightarrow AP-reduction with $S(\epsilon) = \epsilon / \alpha\beta$:

$$\begin{aligned} \frac{cost_A(y)}{OPT_A(x)} &\leq \frac{OPT_A(x) + \beta(cost_B(y') - OPT_B(x'))}{OPT_A(x)} \\ &\leq 1 + \alpha\beta \left(\frac{cost_B(y') - OPT_B(x')}{OPT_B(x')} \right) \\ &= 1 + \alpha\beta \left(\underbrace{\frac{cost_B(y')}{OPT_B(x')}} - 1 \right) \\ &\leq 1 + S(\epsilon) = 1 + \epsilon / \alpha\beta \\ &\leq 1 + \epsilon. \quad \square \end{aligned}$$

- also NP reduction

- most popular reduction type

L-reduction \rightarrow PTAS-reduction, max case: (uncovered)

$$\begin{aligned} \text{cost}_A(y) &= \text{OPT}_A(x) - \underbrace{(\text{OPT}_A(x) - \text{cost}_A(y))}_{\leq \beta \cdot (\text{OPT}_B(x') - \text{cost}_B(y'))} \\ &\geq \text{OPT}_A(x) - \beta (\text{OPT}_B(x') - \text{cost}_B(y')) \end{aligned}$$

$$\begin{aligned} \frac{\text{cost}_A(y)}{\text{OPT}_A(x)} &\geq \frac{\text{OPT}_A(x) - \beta (\text{OPT}_B(x') - \text{cost}_B(y'))}{\text{OPT}_A(x)} \\ &= 1 - \beta \frac{\text{OPT}_B(x') - \text{cost}_B(y')}{\text{OPT}_A(x)} \\ &\geq 1 - \alpha \beta \frac{\text{OPT}_B(x') - \text{cost}_B(y')}{\text{OPT}_B(x')} \quad \left. \begin{array}{l} \text{OPT}_B(x') \leq \alpha \cdot \text{OPT}_A(x) \\ \text{OPT}_A(x) \geq \frac{1}{\alpha} \text{OPT}_B(x') \\ \frac{1}{\text{OPT}_A(x)} \leq \alpha \frac{1}{\text{OPT}_B(x')} \\ -\frac{1}{\text{OPT}_A(x)} \geq -\alpha \frac{1}{\text{OPT}_B(x')} \end{array} \right\} \\ &= 1 - \alpha \beta \left(1 - \underbrace{\frac{\text{cost}_B(y')}{\text{OPT}_B(x')}}_{\geq \frac{1}{1+\delta}} \right) \\ &\geq 1 - \alpha \beta + \frac{\alpha \beta}{1+\delta} \\ &= \frac{1}{1+\varepsilon} \quad \text{when } \delta = \frac{1}{\alpha \beta (1 + \frac{1}{\varepsilon}) - 1} = \frac{\varepsilon}{\alpha \beta (1 + \varepsilon - \frac{\varepsilon}{\alpha \beta})} \end{aligned}$$

$$1+\delta = \frac{\alpha \beta (1 + \frac{1}{\varepsilon})}{\alpha \beta (1 + \frac{1}{\varepsilon}) - 1}$$

$$\frac{1}{1+\delta} = \frac{\alpha \beta (1 + \frac{1}{\varepsilon}) - 1}{\alpha \beta (1 + \frac{1}{\varepsilon})}$$

$$\frac{\alpha \beta}{1+\delta} + 1 = \frac{\alpha \beta (1 + \frac{1}{\varepsilon}) + 1}{1 + \frac{1}{\varepsilon}}$$

$$\frac{\alpha \beta}{1+\delta} + 1 - \alpha \beta = \frac{1}{1 + \frac{1}{\varepsilon}} = \frac{1}{\varepsilon + 1}$$


CLEANER: y' is a $(1 - \varepsilon/\alpha\beta)$ -approximation (c < 1 view)
 $\Rightarrow y$ is a $(1 - \varepsilon)$ -approximation
 [Williamson & Shmoys book, 2010]

APX-complete problems:

Max E3SAT-E5: exactly 3 distinct literals/clause
& exactly 5 occurrences/variable]
[Feige - J.ACM 1998]

Max 3SAT-3: [Papadimitriou & Yannakakis - JCSS 1991]

- usual 3SAT \rightarrow 3SAT-3 reduction:



- not approximation preserving: can now set variable x half true & half false at cost of one violation (can't bound damage)
- fix: connect copies x_1, x_2, \dots, x_k with an expander graph where edge is $x_i = x_j$ ($\neg x_i \vee x_j$, $\neg x_j \vee x_i$)
 - \hookrightarrow bounded degree, k nodes
 - $\hookrightarrow \forall \text{cut } (A, B): \# \text{cross edges} \geq \min\{|A|, |B|\}$(simplification of PY91 construction by Crescenzi 1997)
- \Rightarrow setting x_i 's to majority value won't decrease # satisfied clauses
- \Rightarrow 3SAT- $O(1)$ is APX-hard
 - $\hookrightarrow 29 = 2 \cdot 14 + 1$ using 14-regular expander [Lubotzky, Phillips, Sarnak - Combinatorica 1988]
- then use usual reduction \rightarrow 3SAT-3
 - $O(k)$ violations \Leftarrow k violations]
- \Rightarrow L-reduction **actually a gap reduction, not L reduction** \approx

Independent set, max. degree $\Delta = O(1)$

- any maximal indep. set is Δ -approximation
- strict-reduction from Max 3SAT-3

[Papadimitriou & Yannakakis - JCSS 1991]

- variable gadget \Rightarrow indep. set can't use x_i & \bar{x}_i
- clause gadget $\Rightarrow \leq 1$ point, 0 if not satisfied
- max. degree 4
- 3-regular also APX-complete [Berman & Fujito - TCS 1999]

& [Alimonti & Kann - TCS 2000]


Vertex cover

- greedy algorithm is 2-approximation
- L-reduction from Independent set: do nothing [Papadimitriou & Yannakakis - JCSS 1991]
- vertex cover \Leftrightarrow complement is independent
- OPT_{vc} & OPT_{is} both $\Theta(|V|)$
for bounded-degree graphs $\Rightarrow \Theta(\text{each other})$
- absolute error preserved
- 3-regular OK

Dominating set, max. degree $\Delta = O(1)$

- any minimal dominating set is Δ -approximation
- strict-reduction from Vertex cover:

[Papadimitriou & Yannakakis - JCSS 1991]



- \Rightarrow never need to choose edge node (move $\rightarrow v$)
- 3-regular OK